

# THE SCALAR PERTURBATION SPECTRAL INDEX $n_s$ : *WMAP* SENSITIVITY TO UNRESOLVED POINT SOURCES

K. M. HUFFENBERGER,<sup>1,2,3</sup> H. K. ERIKSEN,<sup>4,5</sup> F. K. HANSEN,<sup>4,5</sup> A. J. BANDAY,<sup>6</sup> K. M. GÓRSKI<sup>2,3,7</sup>

(Dated: Received - / Accepted -)  
 Draft version February 2, 2008

## ABSTRACT

Precision measurement of the scalar perturbation spectral index,  $n_s$ , from the *Wilkinson Microwave Anisotropy Probe* temperature angular power spectrum requires the subtraction of unresolved point source power. Here we reconsider this issue, attempting to resolve inconsistencies found in the literature. First, we note a peculiarity in the *WMAP* temperature likelihood's response to the source correction: Cosmological parameters do not respond to increased source errors. An alternative and more direct method for treating this error term acts more sensibly, and also shifts  $n_s$  by  $\sim 0.3\sigma$  closer to unity. Second, we re-examine the source fit used to correct the power spectrum. This fit depends strongly on the galactic cut and the weighting of the map, indicating that either the source population or masking procedure is not isotropic. Jackknife tests appear inconsistent, causing us to assign large uncertainties to account for possible systematics. Third, we note that the *WMAP* team's spectrum was computed with two different weighting schemes: uniform weights transition to inverse noise variance weights at  $l = 500$ . The fit depends on such weighting schemes, so different corrections apply to each multipole range. For the Kp2 mask used in cosmological analysis, we prefer source corrections  $A = 0.012 \pm 0.005 \mu\text{K}^2$  for uniform weighting and  $A = 0.015 \pm 0.005 \mu\text{K}^2$  for  $N_{\text{obs}}$  weighting. Correcting *WMAP*'s spectrum correspondingly, we compute cosmological parameters with our alternative likelihood, finding  $n_s = 0.970 \pm 0.017$  and  $\sigma_8 = 0.778 \pm 0.045$ . This  $n_s$  is only  $1.8\sigma$  from unity, compared to the  $\sim 2.6\sigma$  *WMAP* 3-year result. Finally, an anomalous feature in the source spectrum at  $l < 200$  remains, most strongly associated with W-band.

*Subject headings:* cosmology: observations, cosmic microwave background, cosmological parameters, methods: data analysis

## 1. INTRODUCTION

Measuring  $n_s$ , the spectral index of initial scalar fluctuations, which is scale invariant ( $n_s = 1$ ) in the Harrison-Zeldovich model and slightly shallower in inflation models, is difficult, primarily because experimental systematics require control over a broad range of spatial scales. In inflation, the deviation from unity closely relates to the inflationary potential and the number of  $e$ -folds of expansion, so a statistically robust measurement of  $n_s \neq 1$  places compelling constraints on the physics of the inflationary epoch.

Because all-sky measurements of the cosmic microwave background (CMB) access the largest observable scales in the universe, the angular power spectrum of the CMB, with a long lever arm, is crucial to such studies. Indeed, the latest data release from the *Wilkinson Microwave Anisotropy Probe* (*WMAP*) claims  $\sim 2.6\sigma$  deviation from the Harrison-Zeldovich spectrum (Spergel et al. 2007). Unfortunately, the CMB is not a totally clean measurement. For example, the well-known degeneracy with

the optical depth since recombination ( $\tau$ ) makes precision measurement of  $n_s$  impossible using CMB temperature anisotropies alone, and polarization is required to break it. Complicated noise properties and hints of unknown systematics in the *WMAP* measurement of large-scale polarization indicate that the systematic uncertainty in both  $\tau$  and  $n_s$  should still be considered significant (Eriksen et al. 2007).

Another important, but under-appreciated, complication for the measurement of  $n_s$  is additional power in the angular spectrum from unresolved, and unmasked, point sources. At high  $l$ , this shot noise can significantly bias the power spectrum, and consequently  $n_s$ . The *WMAP* team devised a sensible prescription for dealing with this contaminant: 1) Use the spectral energy distribution measured from detected sources (and distinct from the CMB) to infer it for undetected ones; 2) measure the contamination using multifrequency data; 3) correct the spectrum; and 4) marginalize over the measurement error when computing the likelihood (Hinshaw et al. 2003, 2007).

Huffenberger et al. (2004) found a level of source contamination consistent with the level in the first *WMAP* data release (Hinshaw et al. 2003). However, based on the three year temperature data (Hinshaw et al. 2007), Huffenberger et al. (2006) measured a point source spectrum with two irregularities. First, at  $l > 200$  the spectrum is white, but with an amplitude below the value in the original preprint of Hinshaw et al. (2007). In the present work, we discovered a small error in the power spectra used for the Huffenberger et al. (2006) estimate,

<sup>1</sup> huffenbe@jpl.nasa.gov

<sup>2</sup> Jet Propulsion Laboratory, 4800 Oak Grove Drive, Pasadena CA 91109

<sup>3</sup> California Institute of Technology, Pasadena, CA 91125

<sup>4</sup> Institute of Theoretical Astrophysics, University of Oslo, P.O. Box 1029 Blindern, N-0315 Oslo, Norway

<sup>5</sup> Centre of Mathematics for Applications, University of Oslo, P.O. Box 1053 Blindern, N-0316 Oslo

<sup>6</sup> Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, Postfach 1317, D-85741 Garching bei München, Germany

<sup>7</sup> Warsaw University Observatory, Aleje Ujazdowskie 4, 00-478 Warszawa, Poland

which should have reported  $A = 0.013 \pm 0.001 \mu\text{K}^2$  instead of  $A = 0.011 \pm 0.001 \mu\text{K}^2$ , still below the original WMAP value of  $A = 0.017 \pm 0.002 \mu\text{K}^2$ . Prompted by our result, Hinshaw et al. (2007) re-examined the issue, revising their value down somewhat and increasing the error bars, to  $A = 0.014 \pm 0.003 \mu\text{K}^2$ . The Spergel et al. (2007) bispectrum analysis indicates a non-Gaussianity consistent with these values, but lacks the statistical power of the multifrequency power spectrum comparison. The second peculiarity is that the power at  $100 < l < 200$  in Hufenberger et al. (2006) was inconsistent, at strong statistical significance, with the rest of the white spectrum.

This paper again considers the power spectrum source correction procedure in detail. We begin in Section 2 with a study of the impact of the source correction on the scalar spectral index through the likelihood. Following this, we probe the source amplitude in Section 3, examining the dependence of the fit on the sky weighting, mask, year of observation, and frequency dependence, and present our best estimates of the cosmological parameters. These same tests probe the robustness of the  $l < 200$  feature. Finally, we conclude in Section 4.

## 2. SOURCE CORRECTION IMPACT ON SPECTRAL INDEX

The final WMAP temperature power spectrum is a noise-weighted combination of cross spectra computed from V- (61 GHz) and W-band (94 GHz) maps. Prior to the computation of the angular spectrum, a foreground model is removed from the maps. The individual cross-spectra are corrected for the sky mask, instrument beams, and point source contamination before combination (see Hinshaw et al. 2007). The combined spectrum is folded into the likelihood calculation,<sup>8</sup> which interfaces to a Markov Chain Monte Carlo code such as *CosmoMC* (Lewis & Bridle 2002), yielding parameter estimates.

Assuming the angular spectrum for sources is white, we wish to explore the dependence of  $n_s$  on the size of the source correction. In Figure 1, we show the correction made by Hinshaw et al. (2007). The correction is not white, because the combined spectrum gets more contribution from W-band at higher  $l$ , due to the noise weighting. To change the amplitude of the correction, we simply scale. Later, we discuss the estimation of this amplitude.

We explore two methods for incorporating the source errors into the likelihood. In the first, we use the standard WMAP likelihood method. The total log-likelihood is split in two,  $-2 \log L = \mathcal{L} = \mathcal{L}_0 + \mathcal{L}_1$ , where  $\mathcal{L}_0$  is due to cosmic variance, noise, and mask-induced mode coupling, and  $\mathcal{L}_1$  includes the uncertainty due to the point source and beam errors (Hinshaw et al. 2003, 2007). The computation of  $\mathcal{L}_1$  assumes that total likelihood is Gaussian, and uses the Woodbury formula to compute an update to the likelihood. Under the Gaussian assumption this split is exact. However, the  $\mathcal{L}_0$  term is not treated as Gaussian, but computed using the Gaussian plus log-normal approximation (Verde et al. 2003).

Treating the source and beam uncertainty separately is very fast (since the uncertainty can be well approximated by a small number of modes), but not really necessary.

<sup>8</sup> We are using version 2.2.2 of the WMAP likelihood, available at <http://lambda.gsfc.nasa.gov/>.




FIG. 1.— The size of the source correction in Hinshaw et al. (2007).

It avoids the inversion of a  $\sim 1000 \times 1000$  matrix per likelihood evaluation, but the low- $l$  part of the code already inverts a  $\sim 2000 \times 2000$  matrix. Performing this inversion and including the beam and source term with the other sources of error in  $\mathcal{L}_0$  is therefore little additional burden.

We take this step in our second method for incorporating the source errors. We have modified the WMAP likelihood code to integrate the beam/point source covariance matrix into the cosmic variance/noise/mask covariance matrix. (This requires inverting the original Fisher matrix, adding the beam/point source term, and inverting back.) Then we feed the new covariance matrix to the Gaussian plus log-normal approximation. Under this procedure, the change in the likelihood due to the inclusion of the beam/point sources term is  $-2.64$ , compared to  $\mathcal{L}_1 = -1.22$  computed with the Woodbury expansion, for the test theory spectrum included with the WMAP likelihood code (where  $\mathcal{L} \sim 3541$ ).

In the top panel of Figure 2, we show the dependence of  $n_s$  on the source correction. We hold the errors fixed at the Hinshaw et al. (2007) value of  $\sigma_A = 0.003 \mu\text{K}^2$ , and marginalize over all the other parameters. As the source correction increases, the power spectrum at high  $l$  is lowered, and  $n_s$  decreases. Completely ignoring the source correction shifts  $n_s$  by  $\sim 0.01$  higher, or about  $0.6\sigma$ . Our alternative likelihood procedure shifts  $n_s$  higher by  $0.005$ , or  $0.3\sigma$ .

Next we study the influence of error in the source correction. At very small and at very large error, we expect the parameter measurement to be independent of the error. For very small source error, the errors in other quantities dominate. For large errors, all modes which could be contaminated by point sources are effectively projected out, and the parameter measurement is again independent. Near  $S/N \sim 1$ , the measurement should undergo a transition, where the measurement error increases and (possibly) the mean value changes. In the bottom panel of Figure 2, we show the dependence of  $n_s$  on the error on the source correction amplitude. The WMAP split likelihood method shows an unexpected result. The size of the point source error apparently has no effect on the measurement of  $n_s$ . Even with the point source error rivaling the size of the acoustic peaks, there




FIG. 2.— (Top) The change in  $n_s$  for given source corrections,  $A$ . We compare the *WMAP* team’s likelihood code to our modified version, which computes the high- $l$  likelihood in a slightly different way. The error in the source correction is fixed at the Hinshaw et al. (2007) value used in Spergel et al. (2007). The thin lines bound the 68% probability interval. The Spergel et al. (2007) values are slightly offset horizontally for visibility, and either ignore (higher  $n_s$ ) or include (lower) a correction for Sunyaev-Zeldovich effect contamination. The other  $n_s$  values ignore SZ. (Bottom) Using the *WMAP* team’s source correction, this plot shows the dependence of  $n_s$  on the point source error,  $\sigma_A$ .

is no effect. The values of the likelihood change, but not the distribution of values in the Markov Chain. This seems to be a clear indication that there is something wrong. The dependence of  $n_s$  on  $A$  in the top panel implies that at least the error bars should increase as  $\sigma_A$  increases.

On the other hand, our modified likelihood shows the expected behavior. As the source correction error is made very large, the errors increase by about 38 percent, and the mean value moves above 1. For this likelihood, the modes not subject to contamination by sources actually prefer  $n_s > 1$ , and a solid measurement of the source contamination is vital to the measurement of  $n_s < 1$ . We also note that the errors on the source measurement do not make much difference, as long as  $\sigma_A < 0.003$ . In the next Section, we examine this measurement in more detail.

### 3. UNRESOLVED POINT SOURCE SPECTRA

#### 3.1. Method

The point source spectrum can be estimated via a linear combination of the individual cross-spectra at several frequencies, a combination which projects out the CMB component. Hufenberger et al. (2004, 2006) examined the unresolved source component in *WMAP* data, using a generalized version of the method for the same task from Hinshaw et al. (2003).

In addition to V- and W-band used for cosmological measurement, the source analysis uses Q-band (41 GHz) because the contaminating sources are much brighter at lower frequencies. There are 276 cross-spectra in these three bands, accounting for all combinations of differencing assemblies per channel, and treating the three years of observation separately. These are combined as

$$A_L = \sum_{L'} \sum_{i \neq j} W_{LL'}^{ij} D_{L'}^{ij} \quad (1)$$

to give the point source amplitude  $A_L$  in a band denoted by  $L$ , where the weight is  $W_L^{ij}$  for the binned cross spectrum estimate  $D_L^{ij}$ , made from maps  $i$  and  $j$ . The map cross spectra  $D_L^{ij}$  include the CMB power spectrum and the contribution from sources. By virtue of being cross-spectra ( $i \neq j$ ), they are noisy but do not have a noise bias. The weights are based upon the frequency independence of the CMB signal (in thermodynamic temperature units), the spectral energy distribution of the sources (measured for bright sources as  $\beta \sim -2.0$ ,  $S \propto \nu^{\beta+2}$  (Hinshaw et al. 2003; Trushkin 2003; Hinshaw et al. 2007)), and the estimated noise covariance in the cross spectrum measurements. (See Hufenberger et al. (2004, 2006) for details.)

The weights obey the constraint

$$\sum_{L'} \sum_{i \neq j} W_{LL'}^{ij} = 0, \quad (2)$$

which means that no CMB will leak into the point source estimate if the maps are properly calibrated and the instrumental beams are perfectly deconvolved from the spectra. If the source spectral energy distribution is correct, the weights also provide an unbiased estimate of the point source power spectrum even if the noise covariance is wrong (although an incorrect noise covariance leads to sub-optimal estimates and incorrect error bars).

The covariance of the source power spectrum estimate is

$$\langle A_L A_{L'} \rangle = (\mathbf{W} \mathbf{\Sigma_D} \mathbf{W})_{LL'}, \quad (3)$$

where  $\mathbf{\Sigma_D}$  is the estimate of the cross spectrum covariance matrix. If the covariance matrix is diagonal in cross-spectra and multipole bin, then the weights are diagonal in multipole. Under this assumption, we plot some example weights in Figure 3.

The expected shot noise angular spectrum of sources in the *WMAP* data is flat in  $C_l$ , so we plot our measured spectra as  $l$  versus  $C_l$ . Throughout we normalize the spectrum at Q band in antenna temperature, which gets a larger signal than V or W.

In the following subsections, we describe several tests of the point source measurement to explore the robustness of these features and their origin. Tests included changing the weights on the map for computing the spectrum, modifying the mask, fitting for the galactic hemispheres separately, fitting year by year, and changing the assumed source spectral energy distribution.




FIG. 3.— The weights for various cross-spectra in the source measurement, using the Kp2 mask, and two different map weighting. The individual weights for the 276 cross-spectra were summed together based on frequency coverage, yielding the 6 combinations shown.

### 3.2. Spectrum at $l > 200$

Whatever the cause of the excess power at  $l < 200$  reported by Huffenberger et al. (2006) may be, it is very likely not relevant for correcting the spectrum at high  $l$ . In this Section, we therefore first concentrate on the correction required for the cosmological analysis, considering only  $l \geq 200$ , and then return to the anomalous low- $l$  feature in Section 3.3.

In a maximum likelihood estimate of the power spectrum, the map is weighted by the pixel-pixel inverse signal plus noise covariance. At small scales this estimate is computationally impractical, and Hinshaw et al. (2007) instead approximate it using two weighting schemes in the computation of the power spectra. In the signal dominated regime at  $l < 500$ , they use spectra where every pixel is weighted evenly; in the noise dominated regime at  $l > 500$ , the maps are inverse noise weighted (i.e. weighted by the number of observations,  $N_{\text{obs}}$ ).

We compute the cross-spectrum using these two schemes and estimate the source contribution, plotted in Figure 4. The flat portion of the spectrum is notably higher in the  $N_{\text{obs}}$  weighted case. Specifically, for a white noise fit including  $l > 200$ ,  $A = 0.015 \pm 0.001 \mu\text{K}^2$  versus  $A = 0.012 \pm 0.001 \mu\text{K}^2$  for the flat weighting. (The single amplitude fits for all combinations we tried are compiled in Table 1.) This leads immediately to a key point: Since the combined spectrum is built from two weighting schemes at different  $l$ 's, the source correction must be different as well. Thus we should have a smaller source correction for  $l < 500$  and a larger correction for  $l > 500$ . The weighting scheme used by Hinshaw et al. (2007) is not specified.

This source level difference is a strong indication that




FIG. 4.— Fits for the unmasked source component, comparing spectra computed from maps using uniform weight outside of the mask with using  $N_{\text{obs}}$  (inverse variance) weighting, plotted as the Q-band amplitude. The lowest  $l$ -bin has no detection and is not shown.

TABLE 1  
FITS FOR SOURCE AMPLITUDE

Mask	Weight	Subset	$A$ ( $\mu\text{K}^2$ , ant. Q)	$\sigma_A$	$\chi^2/\text{dof}$
Diagonal Covariance					
30kp0	flat	all	0.006	0.001	0.25
kp2	flat	VW	0.006	0.004	1.99
kp0	flat	all	0.008	0.001	0.54
kp2	flat	noQ1	0.010	0.001	2.02
kp2	flat	noQ2	0.010	0.001	0.71
kp2	flat	yr12	0.011	0.002	1.02
kp2	flat	yr13	0.011	0.002	1.10
kp2	flat	all (N. hemi.)	0.011	0.001	0.92
kp2	flat	all	0.012	0.001	1.39
kp2	flat	yr22	0.012	0.003	1.16
kp2	flat	yr23	0.012	0.002	0.50
kp2	flat	QW	0.013	0.001	0.70
kp2	flat	all (S. hemi.)	0.013	0.001	0.68
kp2	flat	noW1W2	0.013	0.001	2.48
kp2	flat	noW3W4	0.013	0.001	1.23
kp2	flat	QV	0.014	0.001	1.65
kp2	nobs	all	0.015	0.001	0.69
kp2	flat	yr11	0.017	0.003	0.87
kp2	flat	yr33	0.017	0.003	0.20
Diagonal + beams errors					
kp2	flat	noQ1	0.009	0.001	1.87
kp2	flat	VW	0.010	0.004	2.07
kp2	flat	noQ2	0.010	0.001	0.73
kp2	flat	yr12	0.011	0.002	1.05
kp2	flat	yr13	0.011	0.002	1.10
kp2	flat	yr22	0.011	0.003	1.11
kp2	flat	yr23	0.011	0.002	0.34
kp2	flat	all	0.012	0.001	1.53
kp2	flat	QW	0.013	0.001	0.72
kp2	nobs	all	0.015	0.001	0.79
kp2	flat	QV	0.016	0.001	2.30
kp2	flat	yr33	0.017	0.003	0.25
kp2	flat	yr11	0.019	0.003	0.85

NOTE. — Fits for source power spectrum amplitude, including data at  $l > 200$ .






FIG. 5.— The difference map, Q-ILC, shows two point-like objects near the LMC, left, compared to the Q-band noise rms for the same region, right, based on the number of observations reported by the *WMAP* team.

the source population contaminating the *WMAP* spectrum is not isotropic. One possible scenario is the following: Because source positions are stochastic, and source power strongly favors brighter sources ( $C_l \propto \int dS S^2 dN/dS$ ), brighter sources which, by chance, fell in the best observed regions could boost the spectrum in the  $N_{\text{obs}}$  weighting over uniform weighting. However, the size of this effect is much too small to be a viable explanation. We computed power spectra of Monte Carlo realizations of a noiseless map containing isotropically distributed faint sources, based on differential source counts  $dN/dS \propto S^{-2.3}$  (White & Majumdar 2004; Cleary et al. 2005), normalized to *WMAP* source counts at 1 Jy, setting an upper flux limit to reproduce a reasonable amount of power. Over 1000 simulations, for a bin  $100 < l < 150$ , the rms fluctuation of the difference in power between the uniform and  $N_{\text{obs}}$  spectra is 1.8 percent, and should be smaller at higher  $l$ . Thus the odds are very slight that the change in the power is due to chance alignments of sources with the well observed part of the sky; sources at the appropriate flux level are too numerous. Alternatively, the observed anisotropy in the source population could be either be representative of the real sources, or indicate a problem with the masking procedure, with bright sources slipping through.

The  $N_{\text{obs}}$  weighting emphasizes the ecliptic poles, so these are the best places to look for suspect sources. In particular, visual inspection of a Q-ILC difference map (Figure 5) shows two bright sources near the Large Magellanic Cloud, in the highly observed portion of the sky near the south ecliptic pole. These sources are as bright as some of the sources found by the *WMAP* source-detection algorithm, and indeed the brighter one is included in the subsequent catalogs of López-Caniego et al. (2007) and Nie & Zhang (2007), based on *WMAP* data. The presence of the nearby LMC may have interfered with the source finding procedure.

Changing the sky cut gives further indications of an anisotropic source population. In addition to the Kp2 cut used in the cosmological analysis, we recomputed the point source fit using cross-spectra generated with two additional masks and uniform weighting: the Kp0 mask and a conservative mask consisting of the union of Kp0 with a  $|b| < 30^\circ$  cut galactic cut (see Figure 6). As seen in Table 1, the source power drops significantly as we expand the masks, indicating that *unmasked* sources are




FIG. 6.— (Top) Three masks used in this analysis. The smallest, and most aggressive mask is the *WMAP* Kp2 mask, intermediate is Kp0, and most conservative is the union of Kp0 and a  $|b| < 30^\circ$  cut. (Bottom) The source power spectrum, computed with three different masks. As the mask becomes larger, the power goes down, suggesting a concentration of unmasked source power near the galactic plane.

brighter or more common near the plane. This raises two possibilities. Either some of the sources are galactic in origin, or sources are less efficiently found and masked near the plane.

From a visual inspection of the Q-ILC difference map, we find  $\sim 6$  more bright point-like objects, many near the galactic plane, and all in the southern hemisphere (see Table 2). Many of these objects are also already noted in the the López-Caniego et al. (2007) and Nie & Zhang (2007) catalogs. Computing the power spectrum in hemispheres (with the Kp2 mask), the source amplitudes straddle the value for the whole sky. At  $l < 200$ , the Northern hemisphere has slightly less source power and the the Southern slightly more (See Figure 7).

The *WMAP* source mask, built from a variety of catalogs (Hinshaw et al. 2003, 2007), may not cover the sky evenly, because of differences in sensitivity in different regions. It may be more difficult to identify sources amid large galactic foregrounds. This would be particularly true for flat spectrum sources in lower frequency radio surveys. The *WMAP* source mask has noticeable gaps near the galactic plane, but outside the Kp2 and Kp0 masks.

Two jackknife tests, breaking the data into subgroups, also show some peculiarities. First, we divided up the cross spectra based on the years of observation (Figure 8). Each individual year accounts for 28 cross-spectra, while a pair of years accounts for 64. At  $l > 200$ ,




FIG. 7.— The source power spectrum, computed in hemispheres with the Kp2 mask.

TABLE 2  
SOURCES FOUND IN Q-ILC

RA	Dec.	Gal. lon.	Gal. lat.	ID
00 43 14	-73 16 30	303.7	-44.0	PMNJ0047-7308*, <sup>†</sup>
01 04 11	-72 08 02	301.5	-45.0	PMNJ0059-7210*, <sup>†</sup>
04 51 56	-69 31 10	276.2	-33.7	—
05 20 56	-66 10 24	281.0	-35.6	PMNJ0506-6109*, <sup>†</sup>
06 49 57	-16 56 12	227.9	-8.0	PMNJ0650-1637*
20 50 59	+28 51 35	72.5	-9.7	— <sup>†</sup>
20 52 02	+31 55 18	75.1	-8.0	—
20 57 54	+31 29 08	75.6	-9.2	— <sup>†</sup>

NOTE. — Point-like objects, found visually in a Q-ILC difference map. Marked sources (\*,<sup>†</sup>) were found respectively in the catalog of López-Caniego et al. (2007) and Nie & Zhang (2007).




FIG. 8.— Year by year comparisons of the point source power spectrum.

the fits for individual years are always as greater than for pairs of different years. The largest estimate is the (year 3  $\times$  year 3) fit, at  $A = 0.017 \pm 0.003 \mu\text{K}^2$ , while the smallest is (year 1  $\times$  year 2), at  $0.011 \pm 0.002 \mu\text{K}^2$ . This seems unlikely to be due to chance, and could have a




FIG. 9.— The source power spectrum estimate for combinations of bands.

number of causes. For example, a slight cross-correlation between Q1 and Q2 or between Q and V, introducing a small noise bias in the cross spectra, could have this effect. Removing either Q1 or Q2 from the source estimate, the amplitude drops to  $A = 0.010 \pm 0.001 \mu\text{K}^2$ .

However, this dependence on the Q band could also be due to the shape of the source spectrum. For the second jackknife test, we broke the cross spectra up by band. We considered combinations with Q and V bands only, Q and W only, and V and W only. These spectra are shown in Figure 9. At  $l > 200$ , the V+W combinations are very noisy, and have the lowest source fit,  $A = 0.006 \pm 0.004 \mu\text{K}^2$ , though the  $\chi^2$  is poor. The Q+V combination has the largest source amplitude,  $A = 0.016 \pm 0.001 \mu\text{K}^2$ . This discrepancy could mean that the source spectrum is incorrect: If the true spectrum is steeper than the  $\beta = -2$  we have used, or the spectrum steepens between V and W, one could observe this effect. In this case, the Q+V combination gives the most accurate measurement of the Q band source contamination, but this is not used in the cosmological analysis. To get the correction in V and W bands, we are better off to scale from the lower amplitude, although even this is not completely satisfactory. Using the wrong spectrum for sources means that the shape of the source correction will be somewhat incorrect. We gradually steepened the source spectrum, and re-fit. The Q+V, Q+W, and V+W fits can be brought within  $1\sigma$  by setting  $\beta = -2.5$ . These can be made equal (at  $A = 0.012 \mu\text{K}^2$ ) by setting  $\beta \sim -2.9$ , which compared to  $\beta = -2.0$  would represent a 50 percent smaller amplitude in V and an 80 percent smaller amplitude in W. If true, the source correction should be made correspondingly smaller. (We continue to quote values of  $A$  in Q-band assuming  $\beta = -2.0$  scaling to V and W.)

In the appendix, we discuss the impact of beam errors. These tend to have little impact on the spectrum at  $l > 200$ , since the resulting CMB leakage is large where the CMB is large, at lower  $l$ .

Finally, we must decide by how much to correct the combined *WMAP* spectrum. There is a strong detection of unresolved power in the *WMAP* spectra, but as we have seen through the above tests, our knowledge of the

TABLE 3  
COSMOLOGICAL PARAMETERS

Parameter	<i>WMAP</i>	Source/likelihood corrected
<i>WMAP</i> data only		
$\Omega_b h^2$	$0.0223 \pm 0.0007$	$0.0220 \pm 0.0008$
$\Omega_m$	$0.237 \pm 0.034$	$0.242 \pm 0.037$
$h$	$0.735 \pm 0.032$	$0.731 \pm 0.034$
$\tau$	$0.088 \pm 0.030$	$0.090 \pm 0.030$
$n_s$	$0.951 \pm 0.016$	$0.968 \pm 0.017$
$\sigma_8$	$0.742 \pm 0.051$	$0.780 \pm 0.052$
<i>WMAP</i> + ACBAR + BOOMERANG		
$\Omega_b h^2$	$0.0232 \pm 0.0007$	$0.0224 \pm 0.0007$
$\Omega_m$	$0.233 \pm 0.034$	$0.234 \pm 0.032$
$h$	$0.739 \pm 0.033$	$0.742 \pm 0.032$
$\tau$	$0.088 \pm 0.032$	$0.092 \pm 0.030$
$n_s$	$0.951 \pm 0.016$	$0.970 \pm 0.017$
$\sigma_8$	$0.739 \pm 0.051$	$0.778 \pm 0.045$

NOTE. — Comparison of marginalized parameter results obtained from Table 5 of Spergel et al. (2007) (second column) and after our modifications to the source correction and likelihood (third column).

character of these sources is poor. Our best fit amplitudes including all the data are  $A = 0.012 \mu\text{K}^2$  for flat weighting and  $A = 0.015 \mu\text{K}^2$  for  $N_{\text{obs}}$  weighting. But the spectral effects (from considering V and W without Q) indicate it might be much lower, while individual year fits are much higher. We then are forced to the unsatisfying course of artificially inflating the error bars beyond their nominal statistical values, in order to account for these possibilities. Therefore we set the error on the source estimates at  $\sigma_A = 0.005 \mu\text{K}^2$ .

To obtain our final estimate for cosmological parameters, we correct the spectrum for point sources using the two amplitudes quoted above (uniform weights for  $l < 500$  and  $N_{\text{obs}}$  weights for  $l > 500$ ), and our modified likelihood code. The marginalized parameters from the resulting Markov chains are given in Table 3, both considering *WMAP* data alone and including data from ACBAR (Kuo et al. 2004) and BOOMERanG (Jones et al. 2006; Montroy et al. 2006; Piacentini et al. 2006). Because of the new likelihood and the broadened error bars, this new value of  $n_s = 0.970 \pm 0.017$  is only  $1.8\sigma$  away from 1. Another consequence is that  $\sigma_8$  increases to  $0.778 \pm 0.045$ .

### 3.3. Excess at $l < 200$

We now turn to the  $l < 200$  feature, which is inconsistent with a white noise spectrum. This is present in both the uniform and  $N_{\text{obs}}$  weighted spectra, and in each year and pair of years. Because of its shape and prominence at low  $l$ , we initially considered two specific explanations. The first possibility concerned mis-estimation of the overall multiplicative map calibration of each DA map: If two maps are calibrated differently, the weights  $W_L^{ij}$  in Equation 1 would not cancel the CMB component precisely. Thus, one would observe a leakage from the CMB signal into the point source spectrum, with a signature resembling the CMB power spectrum. A similar effect would be caused by beam uncertainties. In Appendix A, we present the formalism to take these uncertainties into account in the method described in Sec-

tion 3.1, and the results from the corresponding analysis of the *WMAP* data are presented in the bottom Section of Table 1. The conclusion from these computations is that neither calibration nor beam uncertainties can explain this effect.

A third hypothesis is residual galactic foregrounds, which should show through the mask and frequency dependency. As the galactic cut widens from Kp2 to the wide cut (Figure 6), the bin from  $l = 100$ –150 drops about  $0.017 \mu\text{K}^2$ . At the same time the fit for the white noise level drops by  $0.006 \mu\text{K}^2$ . Subtracting off the white noise levels for each, the component in the excess has dropped by about 40 percent. The power in the excess is still significant, even for this broad cut.

Next, we turn to the estimates using only two bands. In this case, we see that the feature is strongly enhanced in the V+W combinations, about the same in the Q+W combination, but clearly diminished in the Q+V combination. The latter appears consistent with the flat source spectrum. This may indicate that the excess is associated with the W band. On the other hand, because the W band spectra tend to carry negative weight in these estimates (Figure 3), this excess would represent a deficit of power in W, which is peculiar. It may indicate an over-subtraction of the galactic foreground template in W band, which could have consequences for the cosmological analysis. At this point, it is difficult to be definitive.

## 4. CONCLUSIONS

We have reanalyzed the source correction procedure for the three-year data release of *WMAP*. First, we considered the impact of this procedure in the *WMAP* likelihood code. Surprisingly, we found that the *WMAP* likelihood does not react to changes in the point source correction error. We have devised a modified likelihood which does respond as expected, although we note that more work is needed to completely validate this approach, as it couples to the important problem of how to approximate a non-Gaussian likelihood with fitting formulas over a wide multipole range. To conclude  $n_s < 1$ , a precision measurement of the source contamination is required. We note that the modes not contaminated favor  $n_s$  consistent with unity.

Second, we found several indications that the unmasked source population in *WMAP* data is anisotropic. This implies that the combined spectrum should be corrected differently in two multipole regions, based on the weighting of the map. Anisotropy in the unmasked sources is unexpected, but can be turned to an advantage: By very carefully masking near the ecliptic poles and galaxy, or employing a wide galactic cut, the point source contamination can be cut substantially. This gain must be weighed against the reduction of the sky area.

We note irregularities in jackknife tests of the source fit, grouped by time of observation or frequency band. This prompted us to adopt large errors on the source fit, to account for systematics beyond the estimate of statistical error which accompanies our measurement. This step is necessary to treat the source correction conservatively. With the modified likelihood and reduced source amplitude from ignoring the excess at  $l < 200$ , these enlarged error bars are responsible for raising our values of  $n_s$  and  $\sigma_8$ .



Finally, the previously noted anomalous  $l < 200$  feature is still present, shows signs of being spatially associated with the galaxy, and is most strongly associated with the W band. On the other hand, it does not appear to be associated with calibration or beam errors. It may represent an over-subtraction of the foreground template in W, although further investigation is warranted. However, the immediate conclusion is that this part of the spectrum should not be used to infer the point source amplitude at higher  $l$ 's.

We thank the *WMAP* team for useful discussions and

for providing additional data. In particular we thank Gary Hinshaw, Michael Nolte, and Lyman Page, who suggested examining the effect of beam uncertainties. HEALPix software (Górski et al. 2005) was used to deriving some results in this paper. We also acknowledge use of the Legacy Archive for Microwave Background Data Analysis (LAMBDA). HKE acknowledges financial support from the Research Council of Norway. This work was partially performed at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with NASA.

## REFERENCES

- Cleary, K. A. et al. 2005, MNRAS, 360, 340, arXiv:astro-ph/0412605  
 Eriksen, H. K., Huey, G., Banday, A. J., Górski, K. M., Jewell, J. B., O'Dwyer, I. J., & Wandelt, B. D. 2007, ApJ, 665, L1, arXiv:0705.3643  
 Górski, K. M., Hivon, E., Banday, A. J., Wandelt, B. D., Hansen, F. K., Reinecke, M., & Bartelmann, M. 2005, ApJ, 622, 759, arXiv:astro-ph/0409513  
 Hinshaw, G. et al. 2007, ApJS, 170, 288, arXiv:astro-ph/0603451  
 ——. 2003, ApJS, 148, 135, arXiv:astro-ph/0302217  
 Huffenberger, K. M., Eriksen, H. K., & Hansen, F. K. 2006, ApJ, 651, L81, arXiv:astro-ph/0606538  
 Huffenberger, K. M., Seljak, U., & Makarov, A. 2004, Phys. Rev. D, 70, 063002, arXiv:astro-ph/0404545  
 Jarosik, N. et al. 2007, ApJS, 170, 263, arXiv:astro-ph/0603452  
 Jones, W. C. et al. 2006, ApJ, 647, 823, arXiv:astro-ph/0507494  
 Kuo, C. L. et al. 2004, ApJ, 600, 32, arXiv:astro-ph/0212289  
 Lewis, A., & Bridle, S. 2002, Phys. Rev. D, 66, 103511, arXiv:astro-ph/0205436  
 López-Caniego, M., González-Nuevo, J., Herranz, D., Massardi, M., Sanz, J. L., De Zotti, G., Toffolatti, L., & Argüeso, F. 2007, ApJS, 170, 108, arXiv:astro-ph/0701473  
 Montroy, T. E. et al. 2006, ApJ, 647, 813, arXiv:astro-ph/0507514  
 Nie, J.-Y., & Zhang, S.-N. 2007, Chinese Journal of Astronomy and Astrophysics, 7, 199, arXiv:astro-ph/0611596  
 Page, L. 2007, personal communication  
 Piacentini, F. et al. 2006, ApJ, 647, 833, arXiv:astro-ph/0507507  
 Spergel, D. N. et al. 2007, ApJS, 170, 377, arXiv:astro-ph/0603449  
 Trushkin, S. A. 2003, Bull. Special Astrophys. Obs., 55, 90, arXiv:astro-ph/0307205  
 Verde, L. et al. 2003, ApJS, 148, 195, arXiv:astro-ph/0302218  
 White, M., & Majumdar, S. 2004, ApJ, 602, 565, arXiv:astro-ph/0308464

## APPENDIX

### LEAKAGE FROM CALIBRATION AND BEAM ERRORS

In the appendix, we estimate the leakage from CMB into the point source estimate. We consider two sources of leakage, the overall calibration of the map and the errors in the beam deconvolution. Each of these can cause the CMB spectrum to have a slightly different amplitude or shape in each of the cross spectra. This means that the CMB will not cancel itself in the weighted linear combination to give the source estimate. This leakage can be accounted for in the cross spectrum covariance. The calibration is a constant function of  $l$ , and the beam error is nearly so over a wide range of scales, so the shape of the leakage term strongly follows the CMB power spectrum, which drops rapidly with  $l$ .

The calibration uncertainty is simpler, so we begin there. In the signal (s) dominated regime, without including point sources, we can model the calibration errors with

$$d_i = (1 + g_i)s. \quad (A1)$$

where the calibration in map  $d_i$  is quantified with dimensionless factors  $g_i$  with  $\langle g_i \rangle = 0$  and  $\langle g_i g_j \rangle = \sigma_g^2 \delta_{ij}$ . Then the estimated cross-spectra are

$$D_l^{ij} = (1 + g_i + g_j + g_i g_j) \hat{C}_l. \quad (A2)$$

Where  $\hat{C}_l$  is the power spectrum of the particular sky realization (although in the actual calculation, we substitute the *WMAP* best-fit spectrum). In the cross spectrum  $i \neq j$ , so this is an unbiased estimator,  $\langle D_l^{ij} \rangle = \hat{C}_l$ . The covariance is

$$\text{Cov}(D_l^{ij}, D_{l'}^{pq}) = \hat{C}_l \hat{C}_{l'} [\sigma_g^2 (\delta_{ip} + \delta_{iq} + \delta_{jp} + \delta_{jq}) + \sigma_g^4 (\delta_{ip} \delta_{jq} + \delta_{iq} \delta_{jp})]. \quad (A3)$$

Essentially, the variance only gets a contribution in terms where the two cross-spectra share a map.

The point source estimate is

$$A_l = \sum_{l'} \sum_{(ij)} W_{ll'}^{ij} D_{l'}^{ij} = \sum_{l'} \hat{C}_{l'} \sum_{(ij)} W_{ll'}^{ij} (g_i + g_j + g_i g_j) \quad (A4)$$

where one term has dropped out because the weights sum to zero. Because we have not included any source contribution, we have  $\langle A_l \rangle = 0$ , where the ensemble average is over the calibration factors. For plotting, its useful to compute the variance due to the calibration factors

$$\langle A_l^2 \rangle = \sum_{l' l''} \hat{C}_{l'} \hat{C}_{l''} \sum_{(ij) (pq)} W_{ll'}^{ij} W_{ll''}^{pq} [\sigma_g^2 (\delta_{ip} + \delta_{iq} + \delta_{jp} + \delta_{jq}) + \sigma_g^4 (\delta_{ip} \delta_{jq} + \delta_{iq} \delta_{jp})] \quad (A5)$$






FIG. 10.— (Left) Beam errors taken from Jarosik et al. (2007). The Q-band errors rise smoothly to  $\sim 0.08$  at  $l = 800$ . (Right) rms leakage of the CMB into the point source measurement, with and without taking the contribution to the beam covariance into account.

A word about binning the spectrum is appropriate here. In practice we bin the spectrum because the large number of cross-spectra (276) and multipoles ( $\sim 800$ ) slows the computation, and the signal-to-noise per multipole is low. We may define a binning matrix  $G_{Li}$  which averages quantities in non-overlapping bands indexed by  $L$ , and compute our estimate from the binned cross spectra  $\sum_i G_{Li} D_l^{ij}$ . The quantity we are now computing is the variance in a bin. Since the weights are also computed in bands, this reduces to replacing the  $l$ s with  $L$ s and replacing each power spectrum term  $C_l$  with  $\sum_i G_{Li} \hat{C}_l$ .

In Jarosik et al. (2007), the calibration uncertainty is quoted as 0.5 percent, so  $\sigma_g = 0.005$ . As written, we are assuming the all maps are uncorrelated, the case if the calibration fluctuations are dominated by noise as expected (Page 2007). In this case, fluctuations in the calibration are fairly small compared to the source estimate. If the calibration is dominated by something else, say foregrounds, then whole bands could be correlated, and the calibration leakage can be substantial compared to the source fit.

In a similar way, we can estimate the beam uncertainties. We begin with

$$d_{lm}^i = \hat{B}_l^i (1 + E_l^i) s_{lm}, \quad (\text{A6})$$

where  $\hat{B}_l^i$  is an unbiased measurement of the beam, and  $E_l^i$  is the (small) fractional difference between the true beam and the measured beam. We assume all beams have independent errors, and define a  $\delta$ -function-like object  $\delta_{ij}^B$ , which is unity when maps  $i$  and  $j$  share a beam, and zero otherwise. With this definition, we may use a Gaussian distribution with  $\langle E_l^i \rangle = 0$  and

$$\langle E_l^i E_l^j \rangle = \Sigma_{ll'}^{B,i} \delta_{ij}^B. \quad (\text{A7})$$

That is, the beam errors are correlated in  $l$  but not between beams. The beam-deconvolved cross spectra are

$$D_l^{ij} = (1 + E_l^i)(1 + E_l^j) \hat{C}_l. \quad (\text{A8})$$

Following the same procedure as with gain calibrations, we find a similar expression for the covariance:

$$\text{Cov}(D_l^{ij}, D_{l'}^{pq}) = \hat{C}_l \left[ \Sigma_{ll'}^{B,i} (\delta_{ip}^B + \delta_{iq}^B) + \Sigma_{ll'}^{B,j} (\delta_{jp}^B + \delta_{jq}^B) + \Sigma_{ll'}^{B,i} \Sigma_{ll'}^{B,j} (\delta_{ip}^B \delta_{jq}^B + \delta_{iq}^B \delta_{jp}^B) \right] \hat{C}_{l'}. \quad (\text{A9})$$

*WMAP* found that the beam uncertainty can be well represented as a small number of orthogonal modes ( $\sim 10$ ) Hinshaw et al. (2003, 2007):

$$\Sigma_{ll'}^{B,i} = \sum_r U_{rl}^i U_{r'l'}^i. \quad (\text{A10})$$

where  $r$  denotes the mode. Here the binning is a little more complicated than for the calibration uncertainty. Instead of simply  $\hat{C}_l$ , the quantities which must be binned are  $\hat{C}_l U_l^i$  (for the first-order terms) and  $\hat{C}_l U_{rl}^i U_{tl}^j$  (for the second order term).

At this time, the beam modes for each *WMAP* differencing assembly are not public. (Only the modes for the combined spectrum are included in the likelihood code.) To approximate the beam modes, we take the beam errors from Jarosik et al. (2007) (shown in Figure 10). We treat each of these as a single mode for the associated beam, correlating all the multipoles in that beam. This should provide a conservative estimate of the beam covariance, though it does not capture all of its properties. With these approximate modes, we can compute the beam variance and the rms CMB leakage for any set of weights. This is plotted in the right panel of Figure 10. For our standard diagonal covariance, the rms beam leakage at first looks like a promising explanation for the  $l < 200$  excess. It has a similar shape, and strong correlations bin-to-bin. However, when we recompute the weights taking the beam covariance into

account, we produce a similar estimate for point sources in a combination which allows very little CMB leakage. Given the estimated size of the rms residual CMB, it is surprising that the source estimates are so similar. Perhaps this is an indication that the beam covariance is overestimated. In any case, the  $l < 200$  is either not due to beams, or the approximation for the beam modes is very poor.